

Complete dietetic dry pet food for adult dogs for the dissolution of struvite stones with urine acidifying properties and low level of magnesium.

## RECOMMENDED FOR & NOT RECOMMENDED FOR

- ✓ Dissolution of sterile struvite stones
- ✓ Dissolution of bacterial-associated struvite stones in combination with appropriate antibiotics
- ✓ Prevention of recurrence of struvite uroliths
- ✓ Management of calcium phosphate uroliths (formation and recurrence)
- ✗ Simultaneous use of urine acidifiers
- ✗ Growth & reproduction



1.5 kg, 3 kg and 12 kg

## KEY BENEFITS



### Controlled pH

Promotes an acidic urine to help reduce urinary struvite stone formation and promote dissolution



### Moderate protein

to help minimise substrate availability for urease-producing bacteria



### Great taste

Highly palatable for long term feeding

## ADDITIONAL BENEFITS & CHARACTERISTICS

### Nutritional management of struvite uroliths

**Controlled pH:** formulated to promote an acidic urine

**Moderate protein:** minimise substrate availability for urease-producing bacteria

### Supports the integrity of the urinary tract mucosa

Sources of glycoaminoglycans (GAGs), omega-3 and antioxidants

## COMPOSITION

Corn, wheat flour, dried poultry protein, rice, pork fat, corn protein meal, digest, dried beet pulp, dried egg, minerals, fish oil.

Urine acidifying substances: calcium sulphate, phosphoric acid.

## KEY NUTRIENT VALUES\*

Moisture	7.5%
Protein	22.0%
Fat	15.0%
Carbohydrate	49.1%
Crude fibre	1.5%
Calcium	0.70%
Phosphorus	0.70%
Sodium	0.20%
Potassium	0.70%
Magnesium	0.08%
Chloride	0.70%
Sulphur	0.3%
Vitamin E	305 IU/kg
Metabolisable energy (ME) <sup>†</sup>	3983 kcal/kg

\* Typical analysis in the final product as fed.

<sup>†</sup> Calculated following NRC 2006 equations.

## FEEDING GUIDELINES

Increased water consumption can help dilute the urine and further decrease the risk of crystal formation. For dissolution of struvite stones, an initial feeding period of 5-12 weeks is recommended. For long-term use, an initial feeding period of up to 6 months is recommended, but the dog should be re-evaluated regularly as indicated by the underlying condition.

## DAILY FEEDING QUANTITY

Body weight (kg)	Daily feeding quantity (g/day)
2.5	65
5	105
10	165
15	215
25	300
35	380
45	445
70	600